



Pet Care Tips for cats

What you'll need to know to keep your companion feline happy and healthy.

Background

Cats were domesticated sometime between 4,000 and 8,000 years ago, in Africa and the Middle East. Small wild cats started hanging out where humans stored their grain. When humans saw cats up close and personal, they began to admire felines for their beauty and grace.

There are many different breeds of cats--from the hairless Sphinx and the fluffy Persian to the silvery spotted Egyptian Mau.

But the most popular felines of all are non-pedigree—that includes brown tabbies, black-and-orange tortoiseshells, all-black cats with long hair, striped cats with white socks and everything in between.

Cost

When you first get your cat, you'll need to spend about \$25 for a litter box, \$10 for a collar, and \$30 for a carrier. Food runs about \$170 a year, plus \$50 annually for toys and treats, \$175 annually for litter and an average of \$150 for veterinary care every year.

Note: Make sure you have all your supplies (see our checklist) before you bring your new pet home.

Basic Care Feeding

- An adult cat should be fed one large or two or three smaller meals each day.

- Kittens from 6 to 12 weeks must eat four times a day.
- Kittens from three to six months need to be fed three times a day. You can either feed specific meals, throwing away any leftover canned food after 30 minutes, or keep dry food available at all times. We recommend a high-quality, brand-name kitten or cat food; avoid generic brands. You will need to provide fresh, clean water at all times, and wash and refill water bowls daily.

Although cat owners of old were told to give their pets a saucer of milk, cats do not easily digest cow's milk, which can cause diarrhea in kittens and cats. Treats are yummy for cats, but don't go overboard. Most packaged treats contain lots of sugar and fat, which can pack on the pounds. Some cats like fresh fruits and vegetables, like broccoli, corn or cantaloupe. You can offer these once in a while.

If your kitten is refusing food or isn't eating enough, try soaking her kitten food in warm water. If that doesn't work, kittens can be fed human baby food for a short time. Use turkey or chicken baby food made for children six months and older. Gradually mix with her regular food.

Grooming

Most cats stays relatively clean and rarely need a bath, but you should brush or comb your cat regularly. Frequent brushing helps keep your cat's coat clean, reduces the amount of shedding and cuts down on the incidence of hairballs.

Handling

To pick up your cat, place one hand behind the front legs and another under the hindquarters. Lift gently. Never pick up a cat by the scruff of the neck or by the front legs.

Housing

Your pet should have her own clean, dry place in your home to sleep and rest. Line your cat's bed with a soft, warm blanket or towel. Be sure to wash the bedding often. Please keep your cat indoors. Cats who are allowed outdoors can contract diseases, get ticks or parasites, become lost or get hit by a car, or get into fights with other free-roaming cats and dogs. Also, cats may prey on native wildlife.

Identification

If allowed outdoors (again, we caution against it!), your cat must wear a safety collar and a license tag. A safety collar with an elastic panel will allow your cat to break loose if the collar gets caught on something. And if your pet is indoors-only, a license tag or an implanted microchip can help insure that your cat is returned if he or she becomes lost.

Litter Box

All indoor cats need a litter box, which should be placed in a quiet, accessible location. A bathroom or utility room is a good place for your cat's box. In a multi-level home, one box per floor is recommended. Avoid moving the box unless absolutely necessary. Then do so slowly, a few inches a day. Keep in mind that cats won't use a messy, smelly litter box, so scoop solid wastes out of the box at least once a day. Dump everything, wash with a mild detergent and refill at least once a week; you can do this less frequently if using clumping litter. Don't use ammonia, deodorants or scents, especially lemon, when cleaning the litter box.

Behavior Information

Play

Cats delight in stalking imaginary prey. The best toys are those that can be made to jump and dance around and look alive. Your cat can safely act out her role as a predator by pouncing on toys instead of people's ankles. Please don't use your hands or fingers as play objects with kittens. This type of rough play may cause biting and scratching behaviors to develop as your kitten matures.

Scratching

Cats need to scratch! When a cat scratches, the old outer nail sheath is pulled off and the sharp, smooth claws underneath are exposed. Cutting your cat's nails every two to three weeks will keep them relatively blunt and less likely to harm the arms of both humans and furniture. Provide your cat with a sturdy scratching post, at least three feet high, which will allow her to stretch completely when scratching. The post should also be stable enough that it won't wobble during use, and should be covered with rough material such as sisal, burlap or tree bark. Many cats also like scratching pads. A sprinkle of catnip once or twice a month will keep your cat interested in her post or pad.

Health

Your cat should see the veterinarian at least once a year for an examination and annual shots, and immediately if she is sick or injured.

Ear Mites

These tiny parasites are a common problem that can be transmitted from cat to cat. If your cat is constantly scratching at his ears or shaking his head, he may be infested with ear mites. You will need to call your vet, as your cat's ears will need to be thoroughly cleaned before medication is dispensed.

Feline Urological Syndrome (FUS)

Both males and females can develop this lower urinary inflammation, also called Feline

Lower Urinary Tract Disease

(FLUTD). Signs of FUS include frequent trips to the litter box, blood in the urine and crying out or straining when urinating. If your male cat looks "constipated," he may have a urethral obstruction and can't urinate. This can be fatal if not treated quickly. Urethral blockages are rare in females. About five percent of cats are affected with FUS. Special diets may help prevent this condition.

Fleas and Ticks

Flea infestation should be taken seriously. These tiny parasites feed off of your pet, transmit tapeworms and irritate the skin.

Carefully check your cat once a week for fleas and ticks. If there are fleas on your cat, there will be fleas in your house. You may need to use flea bombs or premise-control sprays, and be sure to treat all animals in your house. Take care that any sprays, powders or shampoos you use are safe for cats, and that all products are compatible when used together. Cats die every year from improper treatment with flea and tick control products. Please contact your veterinarian for the most effective flea control program for your pet.

Medicines and Poisons

Never give your cat medication that has not been prescribed by a veterinarian. For example, did you know that acetaminophin and aspirin can be FATAL to a cat?! Keep rat poison or other rodenticides away from your cat. If you suspect that your animal has ingested a poisonous substance, call your veterinarian or the Animal Poison Control Center for 24-hour animal poison information at (888) 426- 4435.

Spaying and Neutering

Female cats should be spayed and male cats neutered by six months of age. Neutering a male (removing the testicles) can prevent urine spraying, decrease the urge to escape outside and look for a mate, and reduce fighting between males. Spaying a female (removing the ovaries and uterus) helps prevent breast cancer, which is usually fatal, and pyometra (uterus infection), a very serious problem in older females that must be treated with surgery and intensive medical care. Since cats can breed up to three times per year, it is vital that your female feline be spayed to prevent her from having unwanted litters.

Vaccinations

* Kittens should be vaccinated with a combination vaccine (called a "3 in 1") at 2, 3 and 4 months of age, and then annually. This vaccine protects cats from panleukopenia (also called feline distemper), calicivirus and rhinotracheitis. If you have an unvaccinated cat older than four months of age, he will need a series of two vaccinations given 2 to 3 weeks apart, followed by yearly vaccinations.

* There is a vaccine available for feline leukemia virus (FeLV). This is one of the two immune system viruses

(retroviruses) that infect cats. The other is feline immunodeficiency virus (FIV). There is no vaccine available for FIV. Cats can be infected with either virus for months, even years, without any indication that they are carrying a fatal virus. All cats should be tested for these viruses.

FeLV and FIV can be transmitted at birth from the mother or through the bite of an infected cat.

Neither virus can infect humans. Many outdoor and stray cats and kittens carry this infection. Because of the fatal nature of these diseases, you should not expose cats already living in your home by taking in untested cats or kittens. To be safe, keep your cat indoors—but if your cat does go outside, he should be vaccinated against the feline leukemia virus. Remember, no vaccine is 100-percent effective.

Please note, if your companion cat gets sick because he is not properly vaccinated, the vaccinations should be given after your pet has recovered.

Worms

Kittens

and cats can be infected with several types of worms. The key to treatment is correct diagnosis. This will ensure that the medication is effective against the parasite your pet has. A dewormer that eliminates roundworms, for example, will not kill tapeworms. Your veterinarian can best determine the culprit—and prescribe the appropriate medication.

Cat Supply Checklist

- Premium-brand cat food
- Food dish
- Water bowl
- Interactive toys
- Brush
- Comb
- Safety cat collar with ID tag
- Scratching post or scratching pad
- Litter box
- Litter
- Cat carrier
- Cat bed or box with warm blanket or towel